

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

### **Intrauterine Inflammation and Maternal Exposure to Ambient PM<sub>2.5</sub> during Preconception and Specific Periods of Pregnancy: The Boston Birth Cohort**

Rebecca Massa Nachman, Guangyun Mao, Xingyou Zhang, Xiumei Hong, Zhu Chen, Claire Sampankanpanich Soria, Huan He, Guoying Wang, Deanna Caruso, Colleen Pearson, Shyam Biswal, Barry Zuckerman, Marsha Wills-Karp, and Xiaobin Wang

#### **Table of Contents**

---

**Table S1.** Birth-Related Covariates and Characteristics of Subjects Included in or Excluded from the Analysis

**Table S2.** Associations between Maternal Exposure to Ambient PM<sub>2.5</sub> (Quartile) and Odds of IUI in the Study Population for Four Pregnancy Periods, Adjusting for the Other Three Periods

**Table S3.** Spearman Correlation Coefficients for Exposure during Preconception (Pre), Trimester 1 (T1), Trimester 2 (T2), Trimester 3 (T3), the Whole Pregnancy (WP), and the Last Month (LM)

**Table S4.** Associations between Maternal Exposure to Ambient PM<sub>2.5</sub> (Quartile) and Odds of IUI (Subjects within 10 km of a Monitor)

**Table S5.** Associations between Maternal Exposure to Ambient PM<sub>2.5</sub> (Quartile) and Odds of IUI (Subjects within 4 km of a Monitor)

**Table S6.** Associations between Maternal Exposure to Ambient PM<sub>2.5</sub> (Quartile) and Odds of IUI in the Study Population, Stratified by African American or non-African American

**Table S7.** Associations between Maternal Exposure to Ambient PM<sub>2.5</sub> (Quartile) and Odds of IUI in the Study Population, Stratified by Warm (May-Sept) and Cold (Oct-April) Season

**Table S8.** Associations between Maternal Exposure to Ambient PM<sub>2.5</sub> (Quartile) and Odds of IUI in the Study Population, Stratified by Study Periods: 1999-2005 (high PM<sub>2.5</sub> levels) and 2006-2012 (low PM<sub>2.5</sub> levels).

**Figure S1.** Flow Chart of Subjects Enrolled in the Boston Birth Cohort Included in or Excluded from the Current Analysis

**Figure S2.** Mean Ambient PM<sub>2.5</sub> Exposure over Time in a Subset of 5,059 Mothers in the Boston Birth Cohort from 1999-2012